



Please Fax Completed Form To: 888-898-9113

PATIENT INFORMATION (Complete or Fax Existing Chart)		PRESCRIBER INFORMATION		
Name: DOB:		Prescriber Name:		
Address:		State License:		
City, State, Zip:		NPI #: DEA:		
Phone: Alt. Phone:		Address:		
Email: SS#:				
Gender: M F Weight:(lbs) Ht:			Fax:	
Allergies:		Office Contact:	Phone:	
INSURANCE INFORMA	ATION – OR – Send a copy of the patient	's prescription/insurance	cards (front & back)	
Primary Insurance:		RX Card (PBM):		
City, State, Zip:		BIN: PCN:		
Plan #:		City, State, Zip:		
Group #:		Group #:		
Phone:		Phone:		
CLINICAL INFORMATI	ON			
☐ K51.90 Moderate to Severe Ulcerative Colitis		*If PPD test results are not within 12 months, please perform PPD.		
☐ K50.90 Moderate to Severe Crohn's Disease		Tuberculosis Screening: PPD Test Date:		
☐ M06.9 Rheumatoid Arthritis		Results: Negative		
☐ M45.9 Ankylosing Spondylitis		☐ Positive → ☐ Chest X-Ray Performed Date:		
☐ L40.52 Psoriatic Arthritis		X-Ray Results: ☐ Negative		
☐ L40.0 Plaque Psoriasis		☐ Positive → TB treatment Initiated		
☐ Other:		- Positive > 15 treatment initiated		
Labs:				
☐ CBC q: ☐ CN	MP q:		☐ X-Ray: ☐ Other:	
AVSOLA®				
Prescription type: New	σ start \Box Restart \Box Continued therapy	Total Doses Received:	Date of Last Infusion:	
Medication	Directions		Quantity/Refills	
			Loading dose: 3 doses. No refills.	
Avsola® (infliximab-axxq)		dose: 5mg/kg mg IV at week: 0, 2, 6		
	☐ 3mg/kg mg IV at v	mg/kg mg IV at week: 0, 2, 6		
	☐ Other:		week supply	
	☐ Maintenance dose: (mg/kg)	mg IV every weeks	Refill x 1 year unless noted otherwise. Other:	
PRE-MEDICATIONS			Utiler.	
	mg DO or DIV prior to start of infusion			
☐ Diphenhydraminemg, ☐ PO -or- ☐ IV, prior to start of infusion ☐ Acetaminophen 650 mg PO prior to start of infusion				
☐ Prednisonemg, PO -or- ☐ Methylprednisolone 40 mg IVP -or- ☐ Hydrocortisone 100 mg IVP				
□ Other:				
ANAPHYLACTIC REACTION (AR):				
☐ EpiPen® Auto-injector 0.3 mg (1:1000) Inject IM -or- SubQ to patients who weigh ≥ 66 lbs (≥ 30 kg); may repeat in 3-5 mins x 1 if necessary				
□ EpiPen Jr® Auto-injector 0.15mg (1:2000) Inject IM -or- SubQ to patients who weigh 33 - 66 lbs (15-30 kg): may repeat in 3-5 mins x 1 if necessary				
☐ Diphenhydramine 50mg (1mL) - Give 50 mg slow IVP, administer IM if no IV access; may repeat x 1 after 10 mins, if necessary				
\square Hydrocortisone 100mg - Give 100 mg IVP -or- IM if no IV access				
\square Sodium Chloride 0.9% 500 mL infuse IV at a rate of 30 mL/hr				
□ Other:				

Important Information: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.





Please Fax Completed Form To: 888-898-9113

IV ACCESS				
☐ Start PIV if no IV access available ☐ Maintain current central line access				
MONITORING PARAMETERS				
☐ Obtain vital signs and temperature every 15 mins for the 1st hour, then every 30 mins for the remainder of the infusion				
☐ Observe patient for 30 mins following the infusion				
☐ Instruct patient to report symptoms of chills, fever, headache, sore throat, pain, etc.				
☐ Other:				
CATHETER CARE				
☐ Sodium Chloride 0.9% mL IV before and after each IV access and PRN per protocol				
□ Sodium Chloride 0.9% mL as above AND Heparin 100 Units /mL mL IV flush after second saline flush and PRN				
Drawing sharpes weekly and DDN				
□ Dressing changes weekly and PRN □ Antimicrobial dressing PRN				
☐ May obtain blood from IV access for labs ☐ May use Cathflo 2 mg/2 mL sterile water IVP 2 mL per lumen; May repeat after 2 hours x 1				
STANDARD ORDER FOR SIDE EFFECTS				
☐ Promethazine 25 mg − 1-2 tabs po q 4-6 hrs PRN nausea / vomiting ☐ Diphenhydramine 25 mg - 1 to 2 caps po PRN				
☐ Acetaminophen 325 mg - 2 tabs po q 4-6 hrs PRN HA, myalgia, fever ☐ Diphenhydramine 25 mg -or- 50 mg IV x 1 dose PRN	☐ Diphenhydramine 25 mg -or- 50 mg IV x 1 dose PRN			
☐ Promethazine 25 mg IV/IM x 1 dose PRN nausea / vomiting ☐ Other:				
SIGNATURE				
We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral				
X Date:				
Prescriber Signature				

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

Important Information: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.